



7205 W Center Rd, Suite 200 Omaha, NE 68124
515 N 162nd Ave, Suite 102 Omaha, NE 68118
402-397-6600

POST-INSERTION INSTRUCTIONS FOR MEN

- Your insertion site has been covered with two layers of bandages. The inner layer is a steri-strip, and the outer layer is a waterproof dressing.
- **Do not take tub baths or get into a hot tub or swimming pool for 7 days.** You may shower, but do not remove the bandage or steri-strips for 7 days.
- No major exercises for the incision area. No heavy lifting using the legs for 7 days. This includes running, elliptical, squats, lunges, etc. You can do moderate upper body work and normal walking on a flat surface.
- The sodium bicarbonate in the anesthetic may cause the site to swell for 1-3 days.
- The insertion site may be uncomfortable for up to 2 to 3 weeks. If there is itching or redness you may take Benadryl for relief (50 mg orally every 6 hours). Caution: this can cause drowsiness!
- You may experience bruising, swelling, and/or redness of the insertion site which may last from a few days up to 2 to 3 weeks. If the redness worsens after the first 2-3 days, please contact the office.
- You may notice some pinkish or bloody discoloration of the outer bandage. This is normal.
- If you experience bleeding from the incision, apply firm pressure for 5 minutes.
- Please call if you have any bleeding (not oozing) not relieved with pressure, as this is NOT normal.
- Please call if you have any pus coming out of the insertion site, as this is NOT normal.
- We recommend putting an ice pack on the area where the pellets are located a couple of times for about 20 minutes each time over the next 4 to 5 hours. You can continue this for swelling, if needed. Be sure to place something between the ice pack and your bandages/skin. Do not place ice packs directly on bare skin.

REMINDERS:

- Remember to have your post-insertion blood work done 4 weeks after your FIRST insertion.
- Most men will need re-insertion of their pellets 4-5 months after their initial insertion. If you experience symptoms prior to this, please call the office.
- Please call as soon as symptoms that were relieved from the pellets start to return to make an appointment for your next insertion.

ADDITIONAL INSTRUCTIONS:
